

# **BLANK PAGE**



IS: 3241- 1985

# Indian Standard SPECIFICATION FOR GERANYL ACETATE (First Revision)

UDC 665:572:665:524:22



@ Copyright 1986

INDIAN STANDARDS INSTITUTION
MANAK BHAVAN, 9 BAHADUR SHAH ZAFAR MARG
NEW DELHI 110002

# Indian Standard SPECIFICATION FOR GERANYL ACETATE

# (First Revision)

# Natural and Synthetic Perfumery Materials Sectional Committee, PCDC 18

Chairman

Representing

SHRI SHIAM BEHARI LAL

Gupta & Co (P) Ltd, Delhi

Members

SHRI SUDHIR JAIN ( Alternate to

Shri Shiam Behari Lal )

DR AKHTAR HUSAIN

Central Institue of Medicinal and Aromatic Plants (CSIR), Lucknow

DR M. C. NIGAM ( Alternate )

SHRI S. ANWARULLAH DR C. K. ATAL

DR K. L. DHAR ( Alternate )

Bose Institute, Calcutta

PROF S. C. BHATTACHARYYA CHIEF CHEMIST

Central Revenues Control Laboratory, New Delhi

Regional Research Laboratory (CSIR), Jammu

Maschmeijer Aromatics (India) Ltd, Madras

DEPUTY CHIEF CHEMIST ( Alternate )

SHRI A. K. DAS

Directorate General of Technical Development SHRI S. K. LUTHRA ( Alternate )

DR S. S. GOTHOSKAR

Indian Pharmacopoeia Committee (Ministry of Health and Family Welfare)

SHRI A. K. SHRIVASTAVA ( Alternate ) DR RAJENDRA GUPTA

National Bureau of Plant Genetic Resources, New Delhi

DR M. L. MAHESHWARI ( Alternate )

DR S. K. JAIN Botanical Survey of India, Calcutta

SHRI V. MUDGAL ( Alternate )

DR S. U. KULKARNI Hindustan Lever Ltd, Bombay DR S. KRISHNAMOORTHY ( Alternate )

SHRI G. K. LUTHRIA

Mysore Sugandi Dhoop Factory Ltd, Bombay

SHRI P. R. MALHAN

The Development Commissioner (Small Scale Industries), New Delhi

DR S. K. KAPOOR ( Alternate )

(Continued on page 2)

# © Copyright 1986

#### INDIAN STANDARDS INSTITUTION

This publication is protected under the Indian Copyright Act (XIV of 1957) and reproduction in whole or in part by any means except with written permission of the publisher shall be deemed to be an infringement of copyright under the said Act.

# ( Continued from page 1 )

#### Members

PROF H. H. MATHUR PROF G. K. TRIVEDI ( Alternate )

DR R. K. MATHUR DR B. K. SAIKIA ( Alternate )

SHRI C. K. MEHROTRA

SHRIS. S. CHOPRA ( Alternate ) Dr S. K. Mukherjee

DR N. K. Roy ( Alternate )

SHRIT, S. NARAYANASWAMY

SHRI B. M. NATRAJ

DR K. B. PATIL ( Alternate )

DR S. G. PATNEKAR SHRI SUDHIR JAIN ( Alternate )

DR SHRIDHAR GOPAL PATNEKAR DR ASHOK DIGAMBER WAGH ( Alternate )

SHRI B. K. PATNEY SHRI G. S. RANADE ( Alternate )

SHRI P. B. RAO

SHRI V. VAIDYANATHAN ( Alternate )

DR B. RAVINDRANATH

DR K. N. GURUDUTT ( Alternate )

SHRI S. S. REVADI Karnataka Soaps and Detergents Ltd (Sandalwood Oil Division), Mysore

SHRI B. L. GOPALA KRISHNA ( Alternate ) DR S. C. SETHI

DR A. S. RAO ( Alternate ) SHRI M. P. SHIVA SHRI G. S. SHUKLA

DR S. JAYARAMAN ( Alternate )

DR A. P. SINGH SHRIS, N. MEHRA ( Alternate )

SHRI I. SUNDARESH SHRI Y. G. VAIDYA

SHRI S. B. SULE ( Alternate )
SHRI R V. VAZE

SHRI M. S. SAXENA, Director (P&C)

# Representing

Indian Institute of Technology, Bombay

Regional Research Laboratory (CSIR), Jorhat

Export Inspection Council of India, Calcutta

Indian Agricultural Research Institute, New Delhi

Swastik Household & Industrial Products Ltd. Bombay

Karnataka Soaps and Detergents Ltd (Soaps & Detergent Division ), Bangalore

Perfumes & Flavours Association of India, Bombay

Industrial Perfumes Ltd. Bombay

Richardson Hindustan Ltd, Bombay

The Tata Oil Mills Co Ltd, Bombay

Central Food Technological Research Institute (CSIR), Mysore

National Chemical Laboratory (CSIR), Pune

Forest Research Institute & Colleges, Dehra Dun Directorate of Marketing and Inspection, Faridabad

Camphor & Allied Products Ltd, Bareilly

Basic Chemicals. Pharmaceuticals & Cosmetic Export Promotion Council, Bombay

Naarden (India) Ltd, Bombay

S. H. Kelkar & Co Ltd. Bombay Director General, ISI ( Ex-officio Member )

### Secretary

SHRIMATI VIJAY MALIK Assistant Director ( P&C ), ISI


# Indian Standard

# SPECIFICATION FOR GERANYL ACETATE

# (First Revision)

# O. FOREWORD

- **0.1** This Indian Standard (First Revision) was adopted by the Indian Standards Institution on 25 March 1985, after the draft finalized by Natural and Synthetic Perfumery Materials Sectional Committee had been approved by the Petroleum, Coal and Related Products Division Council.
- **0.2** This standard was first published in 1965. The Sectional Committee responsible for its preparation felt that the standard should be revised with a view to bringing it in line with trade practices in perfumery technology and also to align the quality level of the material currently manufactured and sold in the country which is given here for guidance only.
- **0.3** In this revision, the gas chromatographic method which is being progressively used in the country has been included and the wet method of analysis for determination of ester content has been deleted.
- **0.4** Geranyl acetate ( $C_{12}H_{20}O_2$ ) is the acetate of primary allyl alcohol, geraniol. It is also known as 2-(trans)-3, 7-dimethyl-2, 6 octadienely-lacetate. It is represented by the following structural formula:



### IS: 3241 - 1985

- 0.5 Geranyl acetate is distributed widely in nature and occurs in oils of Daucus carota, Eucalyptus macarthurii, Eucalyptus staigeriana and to a smaller extent in oils of geranium, citronella (Java and Formosa types), palmarosa, neroli, petitgrain and others. However geranyl acetate produced commercially in India is obtained from geraniolex citronella oil which itself is a natural mixture of geraniol and citronellol. Hence, geranyl acetate available at present in India is mainly a mixture of geranyl acetate and citronellyl acetate.
- 0.6 The material is widely used in different perfumery compounds and mainly in conjunction with geraniol, the parent alcohol in rose compositions and for artificial oils of geranium and lavender. It is extensively used for compounding many kinds of perfumes for cosmetics, toiletries and soaps, and in the blending of artificial essential oils and flavours.
- **0.7** In the preparation of this standard, considerable assistance has been derived from the following publications:

EOA No. 11-1975 Standard for Geranyl Acetate. Essential Oil Association of USA, New York.

The Givaudan Index, 1978. Givaudan-Delawanna, Inc., New York.

**0.8** For the purpose of deciding whether a particular requirement of this standard is complied with, the final value, observed or calculated, expressing the result of a test or analysis, shall be rounded off in accordance with IS: 2-1960\*. The number of significant places retained in the rounded off value should be the same as that of the specified value in this standard.

# 1. SCOPE

1.1 This standard prescribes the requirements and the methods of sampling and test for the material commercially known as geranyl acetate obtained from oil of citronella (ex citronella oil).

# 2. TERMINOLOGY

2.1 For the purpose of this standard, definitions given in IS: 6597-1972† shall apply.

# 3. REQUIREMENTS

# 3.1 Description

3.1.1 The material is exclusively manufactured by acetylation of geraniol ex citronella oil (see Type 2 of IS: 1800-1961‡). The acetylated product shall have been carefully fractionated.

<sup>\*</sup>Rules for rounding off numerical values ( revised ).

<sup>†</sup>Glossary of terms relating to natural and synthetic perfumery materials.

<sup>†</sup>Specification for geraniol.

- 3.1.2 The material shall be a clear and colourless liquid, free from sediment, suspended matter and adulterants.
- 3.1.3 The material shall also be tested olfactorily and especially for bynotes and for the presence of adulterants and impurities, if any, as prescribed under 4 and 5 of IS: 2284-1963\*.
- 3.2 Solubility The material shall be soluble in 8 volumes of ethanol (70 percent by volume), when tested as prescribed under 8 of IS: 326-1968†.
- 3.3 The material shall also comply with the requirements given in Table 1.

### 4. PACKING AND MARKING

- 4.1 Packing The material shall be supplied in amber-coloured glass or other air-tight opaque tin-lined or aluminium containers. Suitable galvanized iron containers may also be used.
- 4.1.1 The material shall be protected from light and stored in a cool and dry place.
- 4.2 Marking The material shall be marked with the following information:
  - a) Name of the material;
  - b) Name of the manufacturer and recognized trade-mark, if any:
  - c) Net mass of the material; and
  - d) Batch number.
- 4.2.1 The containers may also be marked with the ISI Certification Mark.

Note — The use of the ISI Certification Mark is governed by the provisions of the Indian Standards Institution (Certification Marks) Act and the Rules and Regulations made thereunder. The ISI Mark on products covered by an Indian Standard conveys the assurance that they have been produced to comply with the requirements of that standard under a well-defined system of inspection, testing and quality control which is devised and supervised by ISI and operated by the producer. ISI marked products are also continuously checked by ISI for conformity to that standard as a further safeguard. Details of conditions, under which a licence for the use of the ISI Certification Mark may be granted to manufacturers or processors, may be obtained from the Indian Standards Institution.

<sup>\*</sup>Method for olfactory assessment of natural and synthetic perfumery materials.

<sup>†</sup>Methods of sampling and test for natural and synthetic perfumery materials (first revision).

TABLE 1 REQUIREMENTS FOR GERANYL ACETATE

( Clauses 3.3 and 6.1)

SL No.	CHARACTERISTIC	REQUIREMENT	METHODS OF TEST, REF TO	
			Indian Stan- dard	Appendix
(1)	(2)	(3)	(4)	(5 <b>)</b>
i)	Odour	Fresh, dry rosepetal like note	IS: 2284-1963*	
ii)	Relative density at 27/27°C (see Note 1)	0·899 2 to 0·904 9	IS: 326 ( Part 3)-1980†	
iii)	Optical rotation	± 1°	IS: 326 ( Part 4)-1980†	
iv)	Refractive index at 27°C (see Note 2)	1: <b>4</b> 52 to 1: <b>4</b> 58	Cl 7 of IS: 326-1968‡	
v)	Acid value, Max	1.0	IS: 326 ( Part 7 )-1980†	
vi)	Gas chromatographic analysis:		<del></del>	Α
	a) Geranyl acetate, percent by mass, Min	69		
	b) Citronellyl acetate, percent by mass, Max	25		

Note 1 — The correction factor for relative density for each degree celsius change in temperature is 0.000 64.

Note 2 — The correction factor for refractive index for each degree celsius change in temperature is 0.00038.

# 5. SAMPLING

5.1 Representative samples of the material, each sample containing not less than 50 ml shall be drawn as prescribed in IS: 326 (Part 1)-1984\*.

#### 6. TEST METHODS

**6.1** Tests shall be carried out as prescribed under **3.1**, **3.2**, **3.3** and the appropriate references specified in col 4 and 5 of Table 1.

<sup>\*</sup>Method for olfactory assessment of natural and synthetic perfumery materials.

<sup>†</sup>Methods of sampling and test for natural and synthetic perfumery materials:

Part 3 Relative density ( second revision ).

Part 4 Determination of optical rotation ( second revision ).

Part 7 Determination of acid value.

<sup>†</sup>Methods of sampling and test for natural and synthetic perfumery materials (first revision).

<sup>\*</sup>Methods of sampling and test for natural and synthetic perfumery materials: Part 1 Sampling (second revision).

6.2 Quality of Reagents — Unless specified otherwise, pure chemicals and distilled water (see IS: 1070-1977\*) shall be employed in tests.

Note — 'Pure chemicals' shall mean chemicals that do not contain impurities which affect the results of analysis.

# APPENDIX A

[ Table 1, Item (vi) ]

# GAS CHROMATOGRAPHIC ANALYSIS OF GERANYL ACETATE

# A-0. GENERAL

- **A-0.1** The chromatographic conditions given here are for guidance only.
- A-0.2 Outline of the Method A sample of the material is injected into the gas chromatograph where it is carried by the carrier gas from one end of the column to the other. During its movement the constituents of the sample undergo distribution at different rates and ultimately get separated from one another. The separated constituents emerge from the end of the column one after another and are detected by suitable means whose response is related to the amount of a specific component leaving the column.

# A-1. APPARATUS

Sample

**A-1.1** Any gas chromatograph capable of being operated under conditions suitable for resolving the individual constituents into distinct peaks may be used. A typical chromatograph for Geranyl acetate using a chromatograph with the following chromatographic conditions is shown in Fig. 1.

Geranyl acetate

•	·
Column	
a) Material	Stainless steel
b) Length	2·438 m
c) OD	<b>7</b> ·95 <b>m</b> m
d) ID	6·35 mm
e) Stationary phase	Carbowax 20 M, 10 percent by mass
f) Solid support	Chromosorb WAW 80-100 mesh

<sup>\*</sup>Method for olfactory assessment of natural and synthetic perfumery materials.

# IS: 3241 - 1985




Fig. 1 Typical Chromatogram of Geranyl Acetate

# A-2. PROCEDURE

A-2.1 Conduct the flow of the carrier gas and inject the sample at inject port where it is vaporized and well mixed with the carrier gas. This is led into the chromatographic column wherein vaporized constituents of the sample are separated out by virtue of their differing interaction with the stationary phase. As the different constituents pass through the detector, they give signals corresponding to the amount of particular constituents leaving the column. The detector signal, on transmission to the recorder, plots the chart. From the specific area under various peaks corresponding to specific constituents, the quantities of different constituents are determined.

NOTE — For the separation to be efficient, it is necessary that the column is maintained at the temperature suggested throughout the time required for the resolution of the constituents.

# A-3. CALCULATION

- A-3.1 Area Measurement (see Note 1) Since normal peaks approximate a triangle the area is measured by multiplying the peak height times the width of half height. The normal peak base is not taken since large deviations may be observed due to tailing or adsorption. This technique is rapid, simple and fairly accurate when peaks are symmetrical and of reasonable width.
- A-3.2 Area Normalization (see Note 2) By normalizing, it meant, calculating the percentage composition by measuring the area of each and dividing the individual areas by total area, for example,

Percentage of 
$$A = \frac{\text{Area of } A}{\text{Total area}} \times 100$$

Note 1 — Other methods of area measurements, namely, Triangulation, Disc Integrator and Electronic Digital Integrator if fixed with GLC machine would be of great advantage.

NOTE 2 — Internal standardization can be used if pure appropriate internal standard is available. This method is known as relative or indirect calibration.

# IS: 3241 - 1985

( Continued from page 2 )

# Synthetic Perfumery Materials Subcommittee, PCDC 18:2

Convener

Representing

SHRI SHIAM BEHARI LAL

Gupta & Co (P) Ltd, Delhi

Members

SHRI SUDIER JAIN ( Alternate to Shri Shiam Behari Lal)

DR P. K. BANERJEE

SHRI A. K. DAS

Hindustan Lever Ltd. Bombay

Directorate General of Technical Development,

New Delhi

SHRIG. D. KELKAR

DR SHRIDHAR GOPAL PATNEKAR DR ASHOK DIGAMBAR WAGH ( Alternate )

SHRI P. B. RAO

SHRI B. HANUMANTHA RAO

SHRI B. K. BALIGA ( Alternate ) SHRIS. S. REVADI

Tata Oil Mills Co Ltd, Bombay

Bush Boake Allen (India) Ltd, Madras

S.H. Kelkar & Co Pvt Ltd, Bombay

Industrial Perfumes Ltd. Bombay

Karnataka Soaps & Detergents Ltd (Sandalwood Oil Division), Mysore

SHRI B. L. GOPALA ( Alternate )

SHRI K. B. SAKSENA SHRI S. C. SAKSENA ( Alternate

DR S. C. SETHE

DR A. S. RAO ( Alternate ) DR A. P. SINGH

SHRIS. N. MEHRA ( Alternate )

India Turpentine and Rosin Co Ltd, Bareilly National Chemical Laboratory (CSIR), Pune

Comphor and Allied Products Ltd, Bareilly



# INDIAN STANDARDS INSTITUTION

# Headquarters:

Manak Bhavan, 9 Behadur Shah Zafar Marg, NEW DELHI 110002

Telephones: 3 31 01 31, 3 31 13 75 Telegrams: Manaksanstha

( Common to all Offices )

# Regional Offices:

Telephona

\*Western : Manakalaya, E9 MIDC, Marol, Andherl ( East ). 6 32 92 95 BOMBAY 400093

†Eastern : 1/14 C. I. T. Scheme VII M, V. I. P. Road, 38 24 99 Maniktola, CALCUTTA 700054

Southern: C. I. T. Campus, MADRAS 600113 41 24 42 Northern: B69 Phase VII, Industrial Focal Point, 8 73 28

S. A. S. NAGAR 160051 (Punjab)

# Branch Offices :

'Pushpak', Nurmohamed Shalkh Marg, Khanpur,	2 63 48 2 63 49
AHMADABAD 380001	12 63 49
'F' Block, Unity Bldg, Narasimharaja Square,	22 48 05

BANGALORE 560002 Gangotri Complex, Bhadbhada Road, T. T. Nagar. 6 27 16 BHOPAL 462003

22E Kalpana Area, BHUBANESHWAR 751014 5 36 27 5-8-56C L. N. Gupta Marg, HYDERABAD 500001 22 10 83 R14 Yudhister Marg, C Scheme, JAIPUR 302005 6 98 32 117/418 B Sarvodaya Nagar, KANPUR 208005 4 72 92

Patliputra Industrial Estate, PATNA 800013 6 23 08
Hantex Bldg ( 2nd Floor ), Rly Station Road, 32 27

TRIVANDRUM 695001
Inspection Office ( With Sale Point ):

Institution of Engineers (India) Building, 1332 Shivaji Nagar, 5 24 35 PUNE 410005

\*Sales Office in Bombay is at Novelty Chambers, Grant Road, 89 65 28 Bombay 400007

†Sales Office in Calcutta is at 5-Chowringhee Approach, P. O. Princep 27 68 00 Street, Calcutta 700072